



Anoka Equine Veterinary Services  
 16445 70TH ST. N.E. ELK RIVER, MN 55330-6522  
 PHONE: (763) 441-3797 FAX: (763) 441-3683  
 AFTER HOURS EMERGENCY NUMBER (320) 274-7119  
 www.anokaequine.com

Kim A. Voller, D.V.M.  
 Kevin M. Voller, D.V.M.  
 J.C. Thieke, D.V.M.  
 Ingrid O. Borkoski, D.V.M.  
 Rick Marion, D.V.M.  
 Tracy A. Turner, D.V.M.  
 MS, DiplACVS  
 Nadine Salomon, V.M.D.  
 David Schwinghamer, D.V.M.

## PLEASURE HORSE

(MINIMAL TRAVELING/SHOWING AND HOUSED IN A CLOSED HERD)

Treatment	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
EEE/WEE /Tetanus Toxoid				X								
West Nile				X								
Rabies								X				
Influenza/Rhino Complex												
Intranasal Strangles												
***Potomac Horse Fever				(X)	X			(X)				

\*\*\*PHF – If given before mid-May, booster again in August. If given after mid-May only vaccinate once.

NOTE: If your horse has never received a particular vaccination before, they will need a second booster in 2 – 4 weeks.

## SHOW/TRAVELING HORSES

(SHOWING/TRAVELING/TRAINING AND/OR LIVING AT A BOARDING BARN)

Treatment	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
EEE/WEE /Tetanus Toxoid				X								
West Nile				X								
Rabies								X				
Influenza/Rhino Complex				X						Y		
Intranasal Strangles				X						Y		
***Potomac Horse Fever				(X)	X			(X)				

\*\*\*PHF – If given before mid-May, booster again in August. If given after mid-May only vaccinate once.

“Y” – Horses in high exposure situations such as boarding or training facilities or horses who show all year around should consider receiving the respiratory vaccine semi-annually.

NOTE: If your horse has never received a particular vaccination before, they will need a booster in 3 – 4 weeks.

### SAMPLE DEWORMING SCHEDULE- ADULT HORSES

<u>February</u>	<u>April</u>	<u>June</u>	<u>August</u>	<u>October</u>	<u>December</u>
moxidectin	pyrantel pamoate	ivermectin	oxibendazole	ivermectin /praziquantal or moxidectin/praziquantal	oxibendazole

**\*\*Consider fecal egg counts 1-2 times per year to help determine if your deworming protocol is effective. Horses who have had fecal tests and are found to be at low risk may be dewormed as few as 2 times per year.**